

PAH Doctor Discussion Guide for Patients on Treatment

If you have been diagnosed with and are being treated for Pulmonary Arterial Hypertension (PAH), it may be time to follow up with your doctor on your current treatment. It's important to monitor your progress and work with your doctor to ensure that you are on the treatment that's right for you.

This guide is designed to help you get the most out of your conversation with your doctor and help you set your treatment goals.

Questions to ask your doctor

Doctor Visits



- Has my condition changed since my last visit?
- What does the next year with PAH look like for me?
- How often should I come in for check-ups?
- Is there anything I can do to prepare for my appointments with you?
- How do I track my progress with PAH?
- What new goals should I set when it comes to improving my PAH symptoms?

Symptoms and Treatment Options



- When it comes to my symptoms of PAH, is there any room for improvement?
- What happens if I don't feel that my current medication is helping my PAH symptoms?
- Are there other medications that may work for me?
- What can I expect when starting a new treatment?

Resources and Support



- Is there a support program that can help patients like me?
- How do I connect with others who can share their PAH journey with me?
- Is there a way for me to offer information to individuals who may have been recently diagnosed with PAH?
- Are there organized events that I can attend to learn more about my PAH?

Living with PAH



- Should I change my diet to help with my PAH symptoms?
- How do I exercise safely with a PAH diagnosis?
- What are some tips to keep in mind when traveling?
- How do I manage household chores without overdoing it?

Please see Important Safety Information on the following pages. For important risk and use information, [click here](#) for the full Prescribing Information, including Boxed Warning and Medication Guide.

Aim Patient Support Program



Aim Nursing Support

- 24/7 hotline provided by the specialty pharmacy
- At-home visits in consultation with your healthcare professional



Aim Access and Financial Support

Co-pay Assistance Program*

- Reduce out-of-pocket costs: up to 100% of co-pays or coinsurance paid directly to the insurer Patient Assistance Program (PAP)[†]
- Eligible individuals can apply to receive medication free of cost for up to one year. Eligibility and criteria may vary

*Patients must have private commercial insurance with a percentage coinsurance or co-pay requirement. Medicare or Medicaid patients are ineligible. Assistance is for one year, after which, patients must re-apply. Patients must notify the program of any change in their insurance status. Patients in certain states may be ineligible. The co-pay assistance program has an annual cap per patient year.

[†]Medicare Part D patients will be enrolled based on calendar year.



myAim Education & Encouragement

Helpful tools and personalized communications

- Educational support and personalized treatment resources through some of the key stages of treatment
- Tips on what to discuss with your doctor and information about other online resources and support

myjourney

Real-life stories of patients like you living with PAH and CTEPH

Learn more at www.adempas-us.com/myjourney

IMPORTANT SAFETY INFORMATION (continued)

Females who are able to get pregnant must use two acceptable forms of birth control, during Adempas[®] (riociguat) treatment and for one month after stopping Adempas.

- If you have had a tubal sterilization, have a progesterone implant, or have an IUD (intrauterine device), these methods can be used alone and no other form of birth control is needed.
- Talk with your doctor or gynecologist (a doctor who specializes in female reproduction) to find out how to prevent pregnancy during treatment with Adempas.
 - **Do not have unprotected sex. Talk to your doctor or pharmacist right away if you have unprotected sex or if you think your birth control has failed. Your doctor may tell you to use emergency birth control.**
 - **Tell your doctor right away if you miss a menstrual period or think you may be pregnant for any reason.**

If you are the parent or caregiver of a female child who started taking Adempas before reaching puberty, you should check your child regularly to see if she is developing signs of puberty. Tell your doctor right away if you notice that she is developing breast buds or any pubic hair. Your doctor should decide if your child has reached puberty. **Your child may reach puberty before having her first menstrual period.**

Females can only receive Adempas through a restricted program called the Adempas Risk Evaluation and Mitigation Strategies (REMS) program. If you are a female who can become pregnant, you must talk to your doctor, understand the benefits and risks of Adempas, and agree to all of the instructions in the Adempas REMS program. Males can receive Adempas without taking part in the Adempas REMS program.

Please see additional Important Safety Information on the following pages. For important risk and use information, [click here](#) for the full Prescribing Information, including Boxed Warning and Medication Guide.



IMPORTANT SAFETY INFORMATION (continued)

Do not take Adempas if:

- **you are pregnant, plan to become pregnant, or become pregnant during treatment with Adempas. Adempas can cause serious birth defects.**
- **you take:**
 - a nitrate medicine to treat high blood pressure or heart disease, such as nitroglycerin, or a medicine called a nitric oxide donor, such as amyl nitrite
 - certain other medicines that contain sildenafil (Revatio or Viagra), tadalafil (Adcirca or Cialis), vardenafil (Levitra or Staxyn), dipyridamole, or theophylline. Revatio and Adcirca are also used to treat PAH
- **you have pulmonary hypertension associated with idiopathic interstitial pneumonias (PH-IIP).**

Ask your doctor or pharmacist if you are not sure if you take any of the medicines listed above.

Before you take Adempas® (riociguat), tell your doctor if you:

- smoke; have recently had serious bleeding from your lungs, or if you have had a medical procedure called bronchial arterial embolization to stop you from coughing up blood; have problems with your heart or blood circulation; have low blood pressure; have liver problems; have kidney problems or are on dialysis; have narrowing of the pulmonary veins, a condition called pulmonary veno-occlusive disease or PVOD; have any other medical conditions.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Adempas and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your doctor.

What should I avoid while taking Adempas?

- **Do not get pregnant** while taking Adempas. If you miss a menstrual period, or think you might be pregnant, call your doctor right away.
- It is not known if Adempas passes into your breast milk. You should not breastfeed if you take Adempas. Talk to your doctor about the best way to feed your baby if you take Adempas.

- Adempas may make you feel dizzy. **Do not drive, operate machinery, or do other activities that require mental alertness or coordination until you know how Adempas affects you.** Talk with your doctor if you are concerned about when it is safe for you to do these activities.
- **Smoking.** Adempas may not work as well if you smoke during treatment. Tell your doctor if you stop smoking or start smoking during treatment with Adempas, because your dose of Adempas may need to be changed.

Adempas can cause serious side effects including:

- **Serious birth defects.**
- **Reduced blood pressure.** This may cause symptoms of low blood pressure, such as lightheadedness, chest pain, and dizziness especially in people who are dehydrated, have a severe blockage of blood flow out of the heart, or have certain other medical problems. Your doctor will check you for any of these problems.
- **Increased risk of bleeding, including bleeding from the respiratory tract.** Tell your doctor right away if you cough up blood during treatment with Adempas.
- **Worsening of symptoms in people with Pulmonary Venous-Occlusive Disease (PVOD).** If you have PVOD, treatment with Adempas may cause a build-up of fluid in your lungs (pulmonary edema) which may result in shortness of breath. Your doctor may tell you to stop taking Adempas and switch you to a different medicine.

The most common side effects of Adempas are:

- Headache; dizziness; indigestion; swelling of your hands, legs, feet, and ankles (peripheral edema); nausea; diarrhea; and vomiting

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Adempas.

For important risk and use information, please see the full Prescribing Information, including Boxed Warning and Medication Guide, [here](#).

You are encouraged to report negative side effects or quality complaints of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

BAYER, the Bayer Cross, Aim, and Adempas are registered trademarks of Bayer.



Bayer
100 Bayer Boulevard, Whippany, NJ 07981 USA
©2020 Bayer
PP-ADE-US-1910-1 December 2020

